

UNIVERSAL DEFORMATION RINGS FOR THE SYMMETRIC GROUP S_4

FRAUKE M. BLEHER AND GIOVANNA LLOSENT

ABSTRACT. Let k be an algebraically closed field of characteristic 2, and let W be the ring of infinite Witt vectors over k . Let S_4 denote the symmetric group on 4 letters. We determine the universal deformation ring $R(S_4, V)$ for every kS_4 -module V which has stable endomorphism ring k and show that $R(S_4, V)$ is isomorphic to either k , or $W[t]/(t^2, 2t)$, or the group ring $W[\mathbb{Z}/2]$. This gives a positive answer in this case to a question raised by the first author and Chinburg whether the universal deformation ring of a representation of a finite group with stable endomorphism ring k is always isomorphic to a subquotient ring of the group ring over W of a defect group of the modular block associated to the representation.

1. INTRODUCTION

Let k be an algebraically closed field of characteristic $p > 0$ and let $W = W(k)$ be the ring of infinite Witt vectors over k . Let G be a finite group, and suppose V is a finitely generated kG -module. If the stable endomorphism ring $\underline{\text{End}}_{kG}(V)$ is one-dimensional over k , it was shown in [6] that V has a universal deformation ring $R(G, V)$. The ring $R(G, V)$ is universal with respect to deformations of V over complete local commutative Noetherian rings with residue field k (see §2). In [6, 3, 4, 5], the isomorphism types of $R(G, V)$ have been determined for V belonging to cyclic blocks, respectively to various tame blocks with dihedral defect groups with one or three isomorphism classes of simple modules. In the present paper, we will consider the case when V belongs to a particular tame block with two isomorphism classes of simple modules. The key tools used to determine the universal deformation rings in all these cases have been results from modular and ordinary representation theory due to Brauer, Erdmann [14], Linckelmann [16, 17], Carlson-Thévenaz [11], and others.

The main motivation for studying universal deformation rings for finite groups is that this case helps understand ring theoretic properties of universal deformation rings for profinite groups Γ . The latter have become an important tool in number theory, in particular if Γ is a profinite Galois group (see e.g. [12], [21, 20], [9] and their references). In [13], de Smit and Lenstra showed that if Γ is an arbitrary profinite group and V is a finite dimensional vector space over k with a continuous Γ -action which has a universal deformation ring $R(\Gamma, V)$, then $R(\Gamma, V)$ is the inverse limit of the universal deformation rings $R(G, V)$ when G runs over all finite discrete quotients of Γ through which the Γ -action on V factors. Thus to answer questions about the ring structure of $R(\Gamma, V)$, it is natural to first consider the case when $\Gamma = G$ is finite.

Suppose now that k has characteristic 2 and that S_4 denotes the symmetric group on 4 letters. In the present paper, we consider the group ring kS_4 which is its own (principal) block and has two isomorphism classes of simple modules. These are represented by the trivial simple module T_0 and a 2-dimensional simple module T_1 which is inflated from the symmetric group S_3 . Since the Sylow 2-subgroups of S_4 are dihedral groups of order 8, kS_4 is of tame representation type. It was shown in [7, 8] that $R(S_4, T_1) \cong W[t]/(t^2, 2t)$ which provides an example of a universal deformation ring which is not a complete intersection. The results in [7, 8] additionally imply that for every finitely generated kS_4 -module V with stable endomorphism ring k , the universal deformation ring $R(S_4, V)$

1991 *Mathematics Subject Classification.* Primary 20C20; Secondary 16G20.

Key words and phrases. Universal deformation rings, stable endomorphism rings, special biserial algebras.

The first author was supported in part by NSF Grants DMS01-39737 and DMS06-51332 and NSA Grant H98230-06-1-0021.

arises from arithmetic in the following sense. By [8, Lemma 3.3 and Thm. 3.7], there are infinitely many real quadratic fields L such that the Galois group $G_{L,\emptyset}$ of the maximal totally unramified extension of L surjects onto S_4 and $R(G_{L,\emptyset}, V) \cong R(S_4, V)$, where V is viewed as a module for $G_{L,\emptyset}$ via inflation. In particular, our main result, Theorem 1.1, determines all the universal deformation rings $R(G_{L,\emptyset}, V)$.

Our main result is as follows, where Ω denotes the syzygy, or Heller, operator (see for example [1, §20]).

Theorem 1.1. *Let V be a finitely generated indecomposable kS_4 -module with stable endomorphism ring k , and let $\mathfrak{C}$ be the component of the stable Auslander-Reiten quiver of kS_4 containing V .*

- i. *Suppose $\mathfrak{C}$ or $\Omega(\mathfrak{C})$ contains T_0 . Then $\mathfrak{C}$ is of type $\mathbb{Z}A_\infty^\infty$, and all modules in $\mathfrak{C} \cup \Omega(\mathfrak{C})$ have stable endomorphism ring equal to k . In this case, $R(S_4, V)$ is isomorphic to $W[\mathbb{Z}/2]$.*
- ii. *Suppose $\mathfrak{C}$ contains T_1 . Then $\mathfrak{C}$ is of type $\mathbb{Z}A_\infty^\infty$, and $\mathfrak{C} = \Omega(\mathfrak{C})$. In this case, V is isomorphic to $\Omega^j(T_1)$ for some integer j , and $R(S_4, V)$ is isomorphic to $W[t]/(t^2, 2t)$.*
- iii. *Suppose $\mathfrak{C}$ contains a uniserial module Y of length 3 with non-isomorphic top and socle. Then $\mathfrak{C}$ is a 3-tube with Y belonging to its boundary, and $\mathfrak{C} = \Omega(\mathfrak{C})$. In this case, V is isomorphic to Y , $\Omega^2(Y)$ or $\Omega^4(Y)$, and $R(S_4, V)$ is isomorphic to k .*

The only components of the stable Auslander-Reiten quiver of kS_4 containing modules with stable endomorphism ring k are the ones in (i) – (iii).

Note that in parts (i) and (iii), $R(S_4, V)$ is a complete intersection, whereas in part (ii) this is not the case. In all three parts, $R(S_4, V)$ is isomorphic to a subquotient ring of WD_8 when D_8 is a dihedral group of order 8. In particular, this gives a positive answer in case of the unique 2-modular block of S_4 to a question raised by the first author and Chinburg in [6, Question 1.1] whether the universal deformation ring of a representation of a finite group with stable endomorphism ring k is always isomorphic to a subquotient ring of the group ring over W of a defect group of the modular block associated to the representation.

The paper is organized as follows. In §2, we give some background on universal deformation rings. In §3, we state the properties of kS_4 we need to prove Theorem 1.1. In particular, we provide the necessary results concerning the indecomposable modules and the stable Auslander-Reiten quiver of kS_4 . In §4–§7, we prove Theorem 1.1.

Part of this paper constitutes the Ph.D. thesis of the second author under the supervision of the first author [18].

2. PRELIMINARIES

Let k be an algebraically closed field of characteristic $p > 0$, let W be the ring of infinite Witt vectors over k and let F be the fraction field of W . Let $\mathcal{C}$ be the category of all complete local commutative Noetherian rings with residue field k . The morphisms in $\mathcal{C}$ are continuous W -algebra homomorphisms which induce the identity map on k .

Suppose G is a finite group and V is a finitely generated kG -module. A lift of V over an object R in $\mathcal{C}$ is a finitely generated RG -module M which is free over R together with a kG -module isomorphism $\phi : k \otimes_R M \rightarrow V$. Two lifts (M, ϕ) and (M', ϕ') of V over R are isomorphic if there is an RG -module isomorphism $\alpha : M \rightarrow M'$ such that $\phi' \circ (1 \otimes \alpha) = \phi$. The isomorphism class of a lift of V over R is called a deformation of V over R , and the set of such deformations is denoted by $\text{Def}_G(V, R)$. The deformation functor $F_V : \mathcal{C} \rightarrow \text{Sets}$ is defined to be the covariant functor which sends an object R in $\mathcal{C}$ to $\text{Def}_G(V, R)$.

Suppose there exists an object $R(G, V)$ in $\mathcal{C}$ and a lift $(U(G, V), \phi_U)$ of V over $R(G, V)$ satisfying the following. For each R in $\mathcal{C}$ and for each lift (M, ϕ) of V over R there is a unique morphism $\alpha : R(G, V) \rightarrow R$ in $\mathcal{C}$ such that the lift (M, ϕ) is isomorphic to $(R \otimes_{R(G, V), \alpha} U(G, V), \phi_U)$ where we identify $k \otimes_R (R \otimes_{R(G, V), \alpha} U(G, V))$ with $k \otimes_{R(G, V)} U(G, V)$. Then $R(G, V)$ is called the universal deformation ring of V and the isomorphism class of the lift $(U(G, V), \phi_U)$ is called the universal deformation of V . In other words, $R(G, V)$ represents the functor F_V in the sense that F_V is

naturally isomorphic to $\text{Hom}_C(R(G, V), -)$. For more information on deformation rings see [13] and [19].

The following four results were proved in [6] and in [4], respectively. As before, Ω denotes the syzygy, or Heller, operator for kG (see for example [1, §20]).

Proposition 2.1. ([6, Prop. 2.1]). *Suppose V is a finitely generated kG -module with stable endomorphism ring $\underline{\text{End}}_{kG}(V) = k$. Then V has a universal deformation ring $R(G, V)$.*

Lemma 2.2. ([6, Cors. 2.5 and 2.8]). *Let V be a finitely generated kG -module with stable endomorphism ring $\underline{\text{End}}_{kG}(V) = k$.*

- i. *Then $\underline{\text{End}}_{kG}(\Omega(V)) = k$, and $R(G, V)$ and $R(G, \Omega(V))$ are isomorphic.*
- ii. *There is a non-projective indecomposable kG -module V_0 (unique up to isomorphism) such that $\underline{\text{End}}_{kG}(V_0) = k$, V is isomorphic to $V_0 \oplus P$ for some projective kG -module P , and $R(G, V)$ and $R(G, V_0)$ are isomorphic.*

Proposition 2.3. ([4, Prop. 2.1.3]). *Let L be a subgroup of G and let U be a finitely generated indecomposable kL -module with $\underline{\text{End}}_{kL}(U) = k$. Suppose there exists an indecomposable kG -module V with $\underline{\text{End}}_{kG}(V) = k$ and a projective kG -module P such that $\text{Ind}_L^G U = V \oplus P$. Assume further that*

$$\dim_k \text{Ext}_{kL}^1(U, U) = \dim_k \text{Ext}_{kG}^1(V, V).$$

Then $R(G, V)$ is isomorphic to $R(L, U)$.

Lemma 2.4. ([4, Lemma 2.3.2]). *Let V be a finitely generated kG -module such that there is a non-split short exact sequence of kG -modules*

$$0 \rightarrow Y_2 \rightarrow V \rightarrow Y_1 \rightarrow 0$$

with $\text{Ext}_{kG}^1(Y_1, Y_2) = k$. Suppose that for $i \in \{1, 2\}$, there exists a WG -module X_i which defines a lift of Y_i over W . Suppose further that

$$\dim_F \text{Hom}_{FG}(F \otimes_W X_1, F \otimes_W X_2) = \dim_k \text{Hom}_{kG}(Y_1, Y_2) - 1.$$

Then there exists a WG -module X which defines a lift of V over W .

3. THE GROUP RING kS_4

Let k be an algebraically closed field of characteristic 2, let W be the ring of infinite Witt vectors over k and let F be the fraction field of W . Then kS_4 is its own principal block, and the defect groups are the Sylow 2-subgroups of S_4 which are dihedral groups of order 8. By [14, §V.2.5.1], kS_4 is Morita equivalent to the special biserial algebra $\Lambda = kQ/I$ where

$$(3.1) \quad Q = \alpha \begin{array}{c} \circlearrowleft \\ \circlearrowright \end{array} \begin{array}{c} 0 \\ \bullet \end{array} \xrightleftharpoons[\gamma]{\beta} \begin{array}{c} 1 \\ \bullet \end{array} \circlearrowright \eta \quad \text{and} \quad I = \langle \alpha^2, \eta\beta, \gamma\eta, \beta\gamma, \gamma\beta\alpha - \alpha\gamma\beta, \eta^2 - \beta\alpha\gamma \rangle.$$

We denote the irreducible Λ -modules by S_0 and S_1 , or, using short-hand, by 0 and 1. Then S_0 corresponds to the trivial simple kS_4 -module T_0 , and S_1 corresponds to the two-dimensional simple kS_4 -module T_1 . The radical series of the projective indecomposable Λ -modules (and hence of the projective indecomposable kS_4 -modules) can be described by the following pictures:

$$(3.2) \quad P_0 = \begin{array}{cc} 0 & 1 \\ 0 & 1 \\ 1 & 0 \\ 0 & \end{array}, \quad P_1 = \begin{array}{cc} 1 & 0 \\ 1 & 0 \\ 0 & 1 \end{array}.$$

The field F is a splitting field for S_4 , and the decomposition matrix of kS_4 has the form

$$(3.3) \quad \begin{array}{c} \varphi_0 \ \varphi_1 \\ \chi_1 \\ \chi_2 \\ \chi_3 \\ \chi_4 \\ \chi_5 \end{array} \begin{bmatrix} 1 & 0 \\ 1 & 0 \\ 1 & 1 \\ 1 & 1 \\ 0 & 1 \end{bmatrix}.$$

Because Λ is a special biserial algebra, all indecomposable non-projective Λ -modules are either string or band modules. Since we will use this description extensively, we now give a brief introduction. For more background material, we refer to [10].

For each arrow $\alpha, \beta, \gamma, \eta$ in Q , we define formal inverses $\alpha^{-1}, \beta^{-1}, \gamma^{-1}, \eta^{-1}$ with starting points $s(\alpha^{-1}) = 0 = s(\gamma^{-1})$ and $s(\beta^{-1}) = 1 = s(\eta^{-1})$ and end points $e(\alpha^{-1}) = 0 = e(\beta^{-1})$ and $e(\gamma^{-1}) = 1 = e(\eta^{-1})$. A word w is a sequence $w_1 \cdots w_n$, where w_i is either an arrow or a formal inverse such that $s(w_i) = e(w_{i+1})$ for $1 \leq i \leq n-1$. Define $s(w) = s(w_n)$, $e(w) = e(w_1)$ and $w^{-1} = w_n^{-1} \cdots w_1^{-1}$. There are also empty words 1_0 and 1_1 of length 0 with $e(1_0) = 0 = s(1_0)$, $e(1_1) = 1 = s(1_1)$ and $(1_0)^{-1} = 1_0$, $(1_1)^{-1} = 1_1$. Denote the set of all words by $\mathcal{W}$, and the set of all non-empty words w with $e(w) = s(w)$ by $\mathcal{W}_r$. Let $J = \{\alpha^2, \eta\beta, \beta\gamma, \gamma\eta, \eta^2, \gamma\beta\alpha, \alpha\gamma\beta, \beta\alpha\gamma\}$.

Definition 3.1. Let $\sim_s$ be the equivalence relation on $\mathcal{W}$ with $w \sim_s w'$ if and only if $w = w'$ or $w^{-1} = w'$. Then strings are representatives $w \in \mathcal{W}$ of the equivalence classes under $\sim_s$ with the following property: Either $w = 1_u$ for $u \in \{0, 1\}$, or $w = w_1 \cdots w_n$ where $w_i \neq w_{i+1}^{-1}$ for $1 \leq i \leq n-1$ and no subword of w or its formal inverse belongs to J .

Let $C = w_1 \cdots w_n$ be a string of length n . Then there exists an indecomposable Λ -module $M(C)$, called the string module $M(C)$ corresponding to the string C , which can be described as follows. There is a k -basis $\{z_0, z_1, \dots, z_n\}$ of $M(C)$ such that the action of Λ on $M(C)$ is given by the following representation $\varphi_C : \Lambda \rightarrow \text{Mat}(n+1, k)$. Let $v(i) = e(w_{i+1})$ for $0 \leq i \leq n-1$ and $v(n) = s(w_n)$. Then for each vertex $u \in \{0, 1\}$ and for each arrow $\xi \in \{\alpha, \beta, \gamma, \eta\}$ in Q

$$\varphi_C(u)(z_i) = \begin{cases} z_i & , \text{ if } v(i) = u \\ 0 & , \text{ else} \end{cases} \quad \text{and} \quad \varphi_C(\xi)(z_i) = \begin{cases} z_{i-1} & , \text{ if } w_i = \xi \\ z_{i+1} & , \text{ if } w_{i+1} = \xi^{-1} \\ 0 & , \text{ else} \end{cases}.$$

We will call φ_C the canonical representation and $\{z_0, z_1, \dots, z_n\}$ the canonical k -basis for $M(C)$ relative to the representative C . Note that $M(C) \cong M(C^{-1})$.

The string modules for the empty words are isomorphic to the simple Λ -modules, namely $M(1_0) \cong S_0$ and $M(1_1) \cong S_1$.

Definition 3.2. Let $w = w_1 \cdots w_n \in \mathcal{W}_r$. Then, for $0 \leq i \leq n-1$, the i -th rotation of w is defined to be the word $\rho_i(w) = w_{i+1} \cdots w_n w_1 \cdots w_i$. Let $\sim_r$ be the equivalence relation on $\mathcal{W}_r$ such that $w \sim_r w'$ if and only if $w = \rho_i(w')$ for some i or $w^{-1} = \rho_j(w')$ for some j . Then bands are representatives $w \in \mathcal{W}_r$ of the equivalence classes under $\sim_r$ with the following property: $w = w_1 \cdots w_n$, $n \geq 1$, with $w_i \neq w_{i+1}^{-1}$ and $w_n \neq w_1^{-1}$, such that w is not a power of a smaller word, and, for all positive integers m , no subword of w^m or its formal inverse belongs to J .

Let $B = w_1 \cdots w_n$ be a band of length n . Then for each integer $m > 0$ and each $\lambda \in k^*$ there exists an indecomposable Λ -module $M(B, \lambda, m)$ which is called the band module corresponding to the band B , λ and m . Note that for all i, j

$$M(B, \lambda, m) \cong M(\rho_i(B), \lambda, m) \cong M(\rho_j(B)^{-1}, \lambda, m).$$

Each component of the stable Auslander-Reiten quiver of Λ consists either entirely of string modules or entirely of band modules. The band modules all lie in 1-tubes. The components consisting of string modules are one 1-tube, one 3-tube and infinitely many non-periodic components of type $\mathbb{Z}A_\infty^\infty$.

The irreducible morphisms between string modules can be described using hooks and cohooks. For our algebra Λ , these are defined as follows. Let $\mathcal{M}$ be the set of maximal directed strings, i.e. $\mathcal{M} = \{\gamma\beta, \beta\alpha, \alpha\gamma, \eta\}$.

Definition 3.3. Let S be a string. We say that S starts on a peak (resp. starts in a deep) if $S = S'C$ (resp. $S = S'C^{-1}$) for some string C in $\mathcal{M}$. Dually, we say that S ends on a peak (resp. ends in a deep) if $S = D^{-1}S''$ (resp. $S = DS''$) for some string D in $\mathcal{M}$.


If S does not start on a peak (resp. does not start in a deep), there is a unique arrow ζ and a unique $M \in \mathcal{M}$ such that $S_h = S\zeta M^{-1}$ (resp. $S_c = S\zeta^{-1}M$) is a string. We say S_h (resp. S_c) is obtained from S by adding a hook (resp. a cohook) on the right side.

Dually, if S does not end on a peak (resp. does not end in a deep), there is a unique arrow ξ and a unique $N \in \mathcal{M}$ such that ${}_hS = M\xi^{-1}S$ (resp. ${}_cS = N^{-1}\xi S$) is a string. We say ${}_hS$ (resp. ${}_cS$) is obtained from S by adding a hook (resp. a cohook) on the left side.

All irreducible morphisms between string modules are either canonical injections $M(S) \rightarrow M(S_h)$, $M(S) \rightarrow M({}_hS)$, or canonical projections $M(S_c) \rightarrow M(S)$, $M({}_cS) \rightarrow M(S)$.

In particular, since none of the projective Λ -modules is uniserial, we get the following result. Suppose S is a string of minimal positive length in a component of the stable Auslander-Reiten quiver of Λ of type $\mathbb{Z}A_\infty$. Then near $M(S)$ the stable Auslander-Reiten component looks as in Figure 3.1.

FIGURE 3.1. The stable Auslander-Reiten component near $M(S)$.



In [15], all homomorphisms between string and band modules have been determined. The following remark describes the homomorphisms between string modules using the canonical bases defined in Definition 3.1.

Remark 3.4. Let $M(S)$ (resp. $M(T)$) be a string module with canonical k -basis $\{x_u\}_{u=0}^m$ (resp. $\{y_v\}_{v=0}^n$) relative to the representative S (resp. T). Suppose C is a string such that

- i. $S \sim_s S'C S''$ with $(S'$ of length 0 or $S' = \hat{S}'\zeta_1$) and $(S''$ of length 0 or $S'' = \zeta_2^{-1}\hat{S}'')$, where $S', \hat{S}', S'', \hat{S}''$ are strings and ζ_1, ζ_2 are arrows in Q ; and
- ii. $T \sim_s T'C T''$ with $(T'$ of length 0 or $T' = \hat{T}'\xi_1^{-1}$) and $(T''$ of length 0 or $T'' = \xi_2\hat{T}'')$, where $T', \hat{T}', T'', \hat{T}''$ are strings and ξ_1, ξ_2 are arrows in Q .

Then there exists a non-zero Λ -module homomorphism $\sigma_C : M(S) \rightarrow M(T)$ which factors through $M(C)$ and which sends each element of $\{x_u\}_{u=0}^m$ either to zero or to an element of $\{y_v\}_{v=0}^n$, according

to the relative position of C in S and T , respectively. If e.g. $S = s_1 s_2 \cdots s_m$, $T = t_1 t_2 \cdots t_n$, and $C = s_{i+1} s_{i+2} \cdots s_{i+\ell} = t_{j+\ell}^{-1} t_{j+\ell-1}^{-1} \cdots t_{j+1}^{-1}$, then

$$\sigma_C(x_{i+t}) = y_{j+\ell-t} \text{ for } 0 \leq t \leq \ell, \text{ and } \sigma_C(x_u) = 0 \text{ for all other } u.$$

Note that there may be several choices of S', S'' (resp. T', T'') in (i) (resp. (ii)). In other words, there may be several k -linearly independent homomorphisms factoring through $M(C)$. By [15], every Λ -module homomorphism $\sigma : M(S) \rightarrow M(T)$ is a k -linear combination of homomorphisms which factor through string modules corresponding to strings C satisfying (i) and (ii).

The next result is an easy consequence of [15] and the symmetric shapes of the radical series of P_0 and P_1 as given in (3.2).

Lemma 3.5. *Let ν be the bijection on the arrows in Q which fixes α and η and interchanges β and γ . Let $S = w_1 w_2 \cdots w_n$ be a string of length $n \geq 1$, and define $\nu(S) = \nu(w_1)^{-1} \nu(w_2)^{-1} \cdots \nu(w_n)^{-1}$. Then $\nu(S)$ is a string, $\dim_k \underline{\text{End}}_\Lambda(M(S)) = \dim_k \underline{\text{End}}_\Lambda(M(\nu(S)))$, and $\dim_k \underline{\text{Ext}}_\Lambda^1(M(S), M(S)) = \dim_k \underline{\text{Ext}}_\Lambda^1(M(\nu(S)), M(\nu(S)))$.*

4. THE STABLE AUSLANDER-REITEN COMPONENTS OF kS_4 CONTAINING T_0 AND $\Omega(T_0)$

In this section, we prove part (i) of Theorem 1.1. Let $\Lambda = kQ/I$ where k , Q and I are as in (3.1), i.e. Λ is Morita equivalent to kS_4 .

Let $\mathfrak{C}_0$ be the component of the stable Auslander-Reiten quiver of Λ containing S_0 , and let U be a Λ -module belonging to $\mathfrak{C}_0 \cup \Omega(\mathfrak{C}_0)$. Using the description of the components of the stable Auslander-Reiten quiver of Λ as in §3, we see that $\mathfrak{C}_0$ is of type $\mathbb{Z}A_\infty^\infty$. Moreover, using hooks and cohooks (see Definition 3.3) we obtain the following. There is an $i \in \mathbb{Z}$ such that $\Omega^i(U)$ is isomorphic to one of the following string modules:

$$(4.1) \quad S_0, \quad A_n = M((\alpha\beta^{-1}\gamma^{-1})^n), \quad B_n = M((\alpha^{-1}\gamma\beta)^n), \quad n \geq 1.$$

Lemma 4.1. *For $n \geq 1$, let V_n be an indecomposable kS_4 -module which, under the Morita equivalence, corresponds to the Λ -module A_n or B_n from (4.1). Then $\underline{\text{End}}_{kS_4}(V_n) \cong k \cong \text{Ext}_{kS_4}^1(V_n, V_n)$.*

Proof. We prove this for V_n corresponding to the Λ -module B_n . The case when V_n corresponds to the Λ -module A_n follows then from Lemma 3.5.

We first show how the results in [2] imply that V_n is endo-trivial, in the sense that the kS_4 -module $\text{End}_k(V_n)$ is isomorphic to a direct sum of the trivial simple kS_4 -module $T_0 = k$ and a projective kS_4 -module. Note that if M and N are endo-trivial kS_4 -modules, then so are $M \otimes_k N$ and $\Omega^i(M)$ for all $i \in \mathbb{Z}$. Consider the almost split sequence ending in T_0

$$(4.2) \quad 0 \longrightarrow \Omega^2(T_0) \longrightarrow V_1 \oplus W_1 \longrightarrow T_0 \longrightarrow 0$$

where, by our assumptions, V_1 corresponds to the Λ -module B_1 , and W_1 corresponds to the Λ -module $\Omega^2(A_1)$. Since $\Omega^{-1}(V_1) \oplus \Omega^{-1}(W_1)$ is isomorphic to the heart $\text{rad}(P_{T_0})/\text{soc}(P_{T_0})$ of the projective indecomposable kS_4 -module P_{T_0} with top T_0 , it follows from [2, Lemma 5.4] that V_1 and W_1 are endo-trivial kS_4 -modules. Given an indecomposable kS_4 -module M of odd k -dimension, it follows from [2, Thm. 3.6 and Cor. 4.7] that

$$(4.3) \quad 0 \longrightarrow \Omega^2(T_0) \otimes_k M \longrightarrow (V_1 \otimes_k M) \oplus (W_1 \otimes_k M) \longrightarrow M \longrightarrow 0$$

is the almost split sequence ending in M modulo projective direct summands. Applying this inductively to $M = V_n$ and using that V_{n+1} is a direct summand of the middle term of the almost split sequence ending in V_n , we obtain that V_n is endo-trivial for all $n \geq 1$.

Since V_n is endo-trivial, it follows that $\underline{\text{End}}_{kS_4}(V_n) \cong k$ and that

$$\text{Ext}_{kS_4}^1(V_n, V_n) \cong H^1(S_4, \text{End}_k(V_n)) \cong H^1(S_4, T_0) \cong \text{Ext}_{kS_4}^1(T_0, T_0) \cong k.$$

□

Lemma 4.2. *For $n \geq 1$, let V_n be an indecomposable kS_4 -module which, under the Morita equivalence, corresponds to the Λ -module A_n or B_n from (4.1). Then $R(S_4, V_n)$ is isomorphic to the group ring $W[\mathbb{Z}/2]$.*

Proof. We prove this for V_n corresponding to the Λ -module A_n , the case of V_n corresponding to B_n being similar.

We use the description in [14, Lemma V.2.5] of the Morita equivalence between kS_4 and $\Lambda = kQ/I$ with Q and I as in (3.1). From this it follows that we can take the arrow α to be $(1+h)e_0$ where $h \in S_4$ is a transposition and e_0 is the idempotent corresponding to the projective indecomposable kS_4 -module P_{T_0} with top T_0 . Let C be the cyclic subgroup of S_4 of order two generated by h . Let T_{00} be the uniserial kS_4 -module with radical series T_0, T_0 . Because of our choice of h , it follows that restricting the action of S_4 on T_{00} to C results in a 2-dimensional kC -module with non-trivial action. Considering a representation of the simple kS_4 -module T_1 , it follows that restricting the action of S_4 on T_1 to C also results in a 2-dimensional kC -module with non-trivial action. Hence

$$(4.4) \quad \text{Res}_C^{S_4} T_{00} \cong kC \cong \text{Res}_C^{S_4} T_1.$$

Let $A_0 = S_0$. For $n \geq 1$, we have a short exact sequence of Λ -modules

$$0 \longrightarrow A_{n-1} \longrightarrow A_n \longrightarrow Y \longrightarrow 0$$

where $Y = M(\alpha\beta^{-1})$. If Z is the kS_4 -module corresponding to Y , then (4.4) implies that

$$\text{Res}_C^{S_4} Z \cong kC \oplus kC.$$

Hence it follows by induction that

$$\text{Res}_C^{S_4} V_n \cong k \oplus (kC)^{2n}$$

where k stands for the trivial simple kC -module. Therefore, $\text{Res}_C^{S_4} V_n$ is a kC -module with stable endomorphism ring k which has universal deformation ring $R(C, \text{Res}_C^{S_4} V_n) \cong W[\mathbb{Z}/2]$. Let $(U_{V_n, C}, \phi_{n, C})$ be a universal lift of $\text{Res}_C^{S_4} V_n$ over $W[\mathbb{Z}/2]$, and let (U_{V_n}, ϕ_n) be a universal lift of V_n over $R(S_4, V_n)$. Then there exists a unique W -algebra homomorphism $f_n : W[\mathbb{Z}/2] \rightarrow R(S_4, V_n)$ in $\mathcal{C}$ such that the lift $(\text{Res}_C^{S_4} U_{V_n}, \text{Res}_C^{S_4} \phi_n)$ is isomorphic to the lift $(R(S_4, V_n) \otimes_{W[\mathbb{Z}/2], f_n} U_{V_n, C}, \phi_{n, C})$.

We next show that f_n is surjective for all $n \geq 1$. For this, we consider all morphisms $\rho : R(S_4, V_n) \rightarrow k[\epsilon]/(\epsilon^2)$. There is a non-split short exact sequence of kS_4 -modules

$$(4.5) \quad 0 \longrightarrow V_n \longrightarrow (P_{T_0})^n \oplus T_{00} \longrightarrow V_n \longrightarrow 0.$$

Hence $E_n = (P_{T_0})^n \oplus T_{00}$ defines a non-trivial lift of V_n over $k[\epsilon]/(\epsilon^2)$. Since

$$\text{Res}_C^{S_4} E_n \cong (kC)^{4n} \oplus kC,$$

$\text{Res}_C^{S_4} E_n$ defines a non-trivial lift of $\text{Res}_C^{S_4} V_n$ over $k[\epsilon]/(\epsilon^2)$. Because $\text{Ext}_{kS_4}^1(V_n, V_n) \cong k$ by Lemma 4.1, this implies that as ρ runs through the morphisms $R(S_4, V_n) \rightarrow k[\epsilon]/(\epsilon^2)$, the composition $\rho \circ f_n$ runs through the morphisms $W[\mathbb{Z}/2] \rightarrow k[\epsilon]/(\epsilon^2)$. Hence f_n is surjective.

We now use Lemma 2.4 to prove that V_n has always two non-isomorphic lifts over W . We first show that the kS_4 -module M corresponding to the Λ -string module $M(\gamma\beta)$ has two non-isomorphic lifts over W . For this we consider the representation $\varphi : S_4 \rightarrow \text{Mat}_4(W)$ which is the natural permutation representation of S_4 , i.e. φ is given by

$$\varphi((1, 2)) = \begin{pmatrix} 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad \varphi((1, 2, 3, 4)) = \begin{pmatrix} 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}.$$

Reducing these matrices modulo 2, results in a representation $\bar{\varphi} : S_4 \rightarrow \text{Mat}_4(k)$ which is the representation of a 4-dimensional indecomposable kS_4 -module corresponding to the Λ -string module $M(\gamma\beta)$. We can order the ordinary irreducible characters of S_4 in (3.3) such that χ_1 is the trivial character, χ_2 is the sign character and the F -character of φ is equal to $\chi_1 + \chi_3$. Twisting φ with

the sign character χ_2 , we obtain a new representation $\varphi' : S_4 \rightarrow \text{Mat}_4(W)$ with F -character $\chi_2 + \chi_4$ which modulo 2 defines a representation of S_4 over k that is also equivalent to $\bar{\varphi}$. This means that we get two non-isomorphic lifts (X, ϕ_X) and $(X', \phi_{X'})$ of M over W such that the F -character of X is $\chi_1 + \chi_3$ and the F -character of X' is $\chi_2 + \chi_4$.

Let $V_0 = T_0$. Then it follows from the decomposition matrix (3.3) that there are two non-isomorphic WS_4 -modules $U_{0,1}^W$ and $U_{0,2}^W$ which define lifts of V_0 over W and whose F -characters are given by χ_1 and χ_2 , respectively.

Let $n \geq 1$. Then $\text{Hom}_{kS_4}(M, V_{n-1}) \cong k^n$, $\text{Ext}_{kS_4}^1(M, V_{n-1}) \cong k$, and we have a short exact sequence of kS_4 -modules

$$(4.6) \quad 0 \longrightarrow V_{n-1} \longrightarrow V_n \longrightarrow M \longrightarrow 0.$$

Assume by induction that there are two non-isomorphic WS_4 -modules $U_{n-1,1}^W$, and $U_{n-1,2}^W$ which define lifts of V_{n-1} over W such that the F -character of $U_{n-1,1}^W$ is

$$(4.7) \quad \left\{ \begin{array}{ll} \chi_1 + s(\chi_1 + \chi_3) + s(\chi_2 + \chi_4) & \text{if } n-1 = 2s \\ \chi_1 + s(\chi_1 + \chi_3) + (s+1)(\chi_2 + \chi_4) & \text{if } n-1 = 2s+1 \end{array} \right\}$$

and the F -character of $U_{n-1,2}^W$ is

$$(4.8) \quad \left\{ \begin{array}{ll} \chi_2 + s(\chi_1 + \chi_3) + s(\chi_2 + \chi_4) & \text{if } n-1 = 2s \\ \chi_2 + (s+1)(\chi_1 + \chi_3) + s(\chi_2 + \chi_4) & \text{if } n-1 = 2s+1 \end{array} \right\}.$$

Let $X^F = F \otimes_W X$, $X'^F = F \otimes_W X'$, $U_{n-1,1}^F = F \otimes_W U_{n-1,1}^W$, and $U_{n-1,2}^F = F \otimes_W U_{n-1,2}^W$. Then if $n-1 = 2s$,

$$\dim_F \text{Hom}_{FS_4}(X'^F, U_{n-1,1}^F) = s + s = n-1,$$

and if $n-1 = 2s+1$,

$$\dim_F \text{Hom}_{FS_4}(X^F, U_{n-1,1}^F) = 1 + s + s = n-1.$$

Similarly, if $n-1 = 2s$ then $\dim_F \text{Hom}_{FS_4}(X^F, U_{n-1,2}^F) = n-1$, and if $n-1 = 2s+1$ then $\dim_F \text{Hom}_{FS_4}(X'^F, U_{n-1,2}^F) = n-1$. Hence by Lemma 2.4, there are two non-isomorphic WS_4 -modules $U_{n,1}^W$ and $U_{n,2}^W$ which define lifts of V_n over W and whose F -characters are as in (4.8) and (4.7) when $n-1$ is replaced by n .

Summarizing, we see that for each $n \geq 1$, we have a surjective morphism $f_n : W[\mathbb{Z}/2] \rightarrow R(S_4, V_n)$ in $\mathcal{C}$ and there are two distinct morphisms $R(S_4, V_n) \rightarrow W$ in $\mathcal{C}$. Hence $\text{Spec}(R(S_4, V_n))$ contains both points of the generic fiber of $\text{Spec}(W[\mathbb{Z}/2])$. Since the Zariski closure of these points is all of $\text{Spec}(W[\mathbb{Z}/2])$, this implies that $R(S_4, V_n)$ must be isomorphic to $W[\mathbb{Z}/2]$. This completes the proof of Lemma 4.2. $\square$


Since by [19, §1.4], $R(S_4, T_0) \cong W[\mathbb{Z}/2]$, part (i) of Theorem 1.1 now follows from Lemmas 4.1, 4.2 and 2.2.

5. THE STABLE AUSLANDER-REITEN COMPONENT OF kS_4 CONTAINING T_1

In this section, we prove part (ii) of Theorem 1.1. As before, let $\Lambda = kQ/I$ where k , Q and I are as in (3.1).

Let $\mathfrak{C}_1$ be the component of the stable Auslander-Reiten quiver of Λ containing S_1 . Using the description of the components of the stable Auslander-Reiten quiver of Λ as in §3, we see that $\mathfrak{C}_1$ is of type $\mathbb{Z}A_\infty^\infty$. Moreover near S_1 , $\mathfrak{C}_1$ looks as in Figure 5.1. where S_{00} is the uniserial Λ -module with radical series S_0, S_0 and S_{0011} is the string module $M(\alpha^{-1}\gamma\eta^{-1})$. Hence $\mathfrak{C}_1 = \Omega(\mathfrak{C}_1)$.

Lemma 5.1. *The only Λ -modules belonging to $\mathfrak{C}_1$ which have stable endomorphism ring k are of the form $\Omega^i(S_1)$ for some $i \in \mathbb{Z}$.*

FIGURE 5.1. The stable Auslander-Reiten component $\mathfrak{C}_1$ near S_1 .

Proof. Let U be a Λ -module belonging to $\mathfrak{C}_1$ which is not in the Ω -orbit of S_1 . Using hooks and cohooks (see Definition 3.3) we obtain the following because of Figure 5.1. There exist integers i and $n \geq 1$ such that $\Omega^i(U)$ is isomorphic to one of the following string modules:

$$\begin{aligned} C_{1,n} &= M \left(\alpha^{-1} (\gamma \eta^{-1} \beta \alpha^{-1} \beta^{-1} \eta \gamma^{-1} \alpha^{-1})^{n-1} \right), \\ C_{2,n} &= M \left(\alpha^{-1} (\gamma \eta^{-1} \beta \alpha^{-1} \beta^{-1} \eta \gamma^{-1} \alpha^{-1})^{n-1} \gamma \eta^{-1} \right), \\ C_{3,n} &= M \left(\alpha^{-1} (\gamma \eta^{-1} \beta \alpha^{-1} \beta^{-1} \eta \gamma^{-1} \alpha^{-1})^{n-1} \gamma \eta^{-1} \beta \alpha^{-1} \beta^{-1} \right). \end{aligned}$$

Let $\{x_r^{\ell,n}\}_{r=0}^{n_\ell}$ be the canonical k -basis for $C_{\ell,n}$, $\ell \in \{1, 2, 3\}$. Then the Λ -module endomorphism of $C_{\ell,n}$ which sends $x_0^{\ell,n}$ to $x_1^{\ell,n}$ and all other $x_r^{\ell,n}$ to zero does not factor through a projective Λ -module. $\square$

Since it was proved in [8, Thm. 2.3] that $R(S_4, T_1) \cong W[t]/(t^2, 2t)$, part (ii) of Theorem 1.1 now follows from Lemmas 5.1 and 2.2.

6. THE STABLE AUSLANDER-REITEN COMPONENT OF kS_4 WHICH IS A 3-TUBE

In this section, we prove part (iii) of Theorem 1.1. As before, let $\Lambda = kQ/I$ where k , Q and I are as in (3.1).

Let $\mathfrak{C}$ be a component of the stable Auslander-Reiten quiver of Λ containing a uniserial module X of length 3 with non-isomorphic top and socle. Then X is either X_1 with descending radical series S_0, S_0, S_1 or X_2 with descending radical series S_1, S_0, S_0 . In particular, $X_2 = \Omega(X_1) = \Omega^{-2}(X_1)$, which implies that $\mathfrak{C}$ is the unique 3-tube of the stable Auslander-Reiten quiver of Λ with X_1 and X_2 belonging to its boundary, and $\mathfrak{C} = \Omega(\mathfrak{C})$. The third Λ -module at the boundary of $\mathfrak{C}$ is $\Omega^2(X_1)$ which is the uniserial Λ -module with radical series S_1, S_1 .

Lemma 6.1. *The only Λ -modules belonging to $\mathfrak{C}$ which have stable endomorphism ring k are X_1 , $X_2 = \Omega(X_1)$ and $\Omega^2(X_1)$. Moreover, if Y is a uniserial kS_4 -module of length 3 with non-isomorphic top and socle and $V \in \{Y, \Omega^2(Y), \Omega^4(Y)\}$, then $R(S_4, V) \cong k$.*

Proof. Let U be a Λ -module belonging to $\mathfrak{C}$ but not to its boundary. Using hooks and cohooks (see Definition 3.3) it follows that there exist integers $n \geq 1$ and $i \in \{0, 1, 2\}$ such that $\Omega^i(U)$ is

isomorphic to one of the following string modules:

$$\begin{aligned} X_{-1,n} &= M \left((\gamma^{-1}\alpha^{-1}\gamma\eta^{-1}\beta\alpha^{-1}\beta^{-1}\eta)^{n-1} \gamma^{-1}\alpha^{-1}\gamma\eta^{-1} \right), \\ X_{0,n} &= M \left((\gamma^{-1}\alpha^{-1}\gamma\eta^{-1}\beta\alpha^{-1}\beta^{-1}\eta)^{n-1} \gamma^{-1}\alpha^{-1}\gamma\eta^{-1}\beta\alpha^{-1}\beta^{-1} \right), \\ X_{1,n} &= M \left((\gamma^{-1}\alpha^{-1}\gamma\eta^{-1}\beta\alpha^{-1}\beta^{-1}\eta)^n \gamma^{-1}\alpha^{-1} \right). \end{aligned}$$

Let $\{x_r^{\ell,n}\}_{r=0}^{8n-1+3\ell}$ be the canonical k -basis for $X_{\ell,n}$, $\ell \in \{-1, 0, 1\}$. Then the Λ -module endomorphism of $X_{\ell,n}$ which sends $x_0^{\ell,n}$ to $x_4^{\ell,n}$, and all other $x_r^{\ell,n}$ to zero does not factor through a projective Λ -module. Hence the stable endomorphism ring of $X_{\ell,n}$ has k -dimension at least 2.

By Lemma 2.2, to prove $R(S_4, V) \cong k$, for V as in the statement of the lemma, it is sufficient to consider the case when $V = T_{11}$ is the uniserial kS_4 -module with radical series T_1, T_1 . Let A_4 be the alternating group of order 12 inside S_4 . There are three one-dimensional simple kA_4 -modules E_0, E_1 and E_2 where E_0 is the trivial simple kA_4 -module. Consider the uniserial kA_4 -module E_{12} with descending radical series E_1, E_2 . Then E_{12} lies at the boundary of a 3-tube of the stable Auslander-Reiten quiver of kA_4 , and there is a non-split short exact sequence of kA_4 -modules

$$0 \longrightarrow E_2 \longrightarrow E_{12} \longrightarrow E_1 \longrightarrow 0.$$

By [1, Lemma 8.6(5)], since $\text{Ind}_{A_4}^{S_4} E_1 \cong T_1 \cong \text{Ind}_{A_4}^{S_4} E_2$, there is a non-split short exact sequence of kS_4 -modules

$$0 \longrightarrow T_1 \longrightarrow \text{Ind}_{A_4}^{S_4} E_{12} \longrightarrow T_1 \longrightarrow 0$$

which implies that $T_{11} \cong \text{Ind}_{A_4}^{S_4} E_{12}$. By [3, Prop. 3.4], $\underline{\text{End}}_{kA_4}(E_{12}) = k$ and $R(A_4, E_{12}) \cong k$. Since $\underline{\text{End}}_{kS_4}(T_{11}) = k$ and

$$\text{Ext}_{kS_4}^1(T_{11}, T_{11}) = 0 = \text{Ext}_{kA_4}^1(E_{12}, E_{12}),$$

Proposition 2.3 implies that $R(S_4, T_{11}) \cong R(A_4, E_{12}) \cong k$. $\square$

7. STABLE ENDOMORPHISM RINGS

In this section, we complete the proof of Theorem 1.1, by determining which components of the stable Auslander-Reiten quiver of kS_4 contain modules with stable endomorphism ring k . Recall that kS_4 is Morita equivalent to the special biserial algebra $\Lambda = kQ/I$ where k, Q and I are as in (3.1).

We first consider all components of the stable Auslander-Reiten quiver of Λ of type $\mathbb{Z}A_\infty^\infty$ and determine which of these components contain a module with stable endomorphism ring k .

Proposition 7.1. *Let $\Lambda = kQ/I$ where Q and I are as in §3. Then the components of the stable Auslander-Reiten quiver of Λ of type $\mathbb{Z}A_\infty^\infty$ containing a module with stable endomorphism ring k are precisely the components containing $S_0, \Omega(S_0)$ or S_1 .*

Proof. Let $\mathfrak{C}$ be a component of type $\mathbb{Z}A_\infty^\infty$ of the stable Auslander-Reiten quiver of Λ such that $\mathfrak{C} \cup \Omega(\mathfrak{C})$ contains no simple Λ -module.

Let $M(S)$ be a string module of minimal length in $\mathfrak{C}$. In particular, S cannot start or end in a peak (resp. in a deep), and near $M(S)$ the component $\mathfrak{C}$ looks as in Figure 3.1. Hence it suffices to show that each of the Λ -modules $M(S)$, $M(S_{h\dots h})$ and $M(S_{c\dots c})$ has an endomorphism which does not factor through a projective Λ -module and which is not a scalar multiple of the identity homomorphism.

Using Lemma 3.5 and that S cannot start or end in a peak (resp. in a deep), it follows that one only needs to consider $M(S)$ where

$$(7.1) \quad S = \alpha^{-1}\gamma \dots, \quad S = \gamma^{-1}\alpha \dots, \quad \text{or} \quad S = \beta^{-1}\eta\gamma^{-1} \dots.$$

Note that $M(\alpha^{-1})$ lies in the same stable Auslander-Reiten component as S_1 , and hence has already been considered in Lemma 5.1.

One now goes through all the cases for S in (7.1) and determines suitable endomorphisms that do not factor through a projective Λ -module. We provide a few examples. Let $\{z_r\}_{r=0}^{\ell(S)}$ be the canonical k -basis of the string module $M(S)$ relative to the chosen representative S .

If $S = \alpha^{-1}\gamma \dots$ then the endomorphism sending z_0 to z_1 and all other z_r to zero does not factor through a projective Λ -module, and the same is true for $M(S_{h\dots h})$ and $M(S_{c\dots c})$.

If $S = \gamma^{-1}\alpha \dots$, one needs to consider two possibilities, namely $S = \gamma^{-1}\alpha\beta^{-1} \dots$ and $S = \gamma^{-1}\alpha\gamma \dots$. If $S = \beta^{-1}\eta\gamma^{-1} \dots$, one needs to consider three possibilities, namely $S = \beta^{-1}\eta\gamma^{-1}$, $S = \beta^{-1}\eta\gamma^{-1}\alpha \dots$ and $S = \beta^{-1}\eta\gamma^{-1}\alpha^{-1} \dots$. For example, if $S = \beta^{-1}\eta\gamma^{-1}\alpha \dots$ then the endomorphism sending z_0 to z_3 and all other z_r to zero does not factor through a projective Λ -module, and the same is true for $M(S_{h\dots h})$ and $M(S_{c\dots c})$. $\square$

Next, we consider all components of the stable Auslander-Reiten quiver which are one-tubes. Except for one 1-tube consisting of string modules, all the other 1-tubes consist of band modules. It follows from [15] that if B is a band, $\lambda \in k - \{0\}$ and $n \geq 2$ is an integer, then $\underline{\text{End}}_\Lambda(M(B, \lambda, n))$ has k -dimension at least 2. Hence we can concentrate on the band modules $M(B, \lambda, 1)$. We need the following two definitions.

Definition 7.2. Let B be a band for Λ , $\lambda \in k - \{0\}$ and let $M_{B,\lambda} = M(B, \lambda, 1)$. Suppose S is a string such that

- i. $B \sim_r ST_1$ with $T_1 = \xi_1^{-1}T'_1\xi_2$, where T_1, T'_1 are strings and ξ_1, ξ_2 are arrows in Q ; and
- ii. $B \sim_r ST_2$ with $T_2 = \zeta_1T'_2\zeta_2^{-1}$, where T_2, T'_2 are strings and ζ_1, ζ_2 are arrows in Q .

Then by [15] there exists a non-zero endomorphism of $M_{B,\lambda}$ which factors through $M(S)$. We will call such an endomorphism to be of string type S . Note that there may be several choices of T_1 (resp. T_2) in (i) (resp. (ii)). In other words, there may be several k -linearly independent endomorphisms of string type S . By [15], every endomorphism of $M_{B,\lambda}$ is a k -linear combination of the identity homomorphism and of endomorphisms of string type S for suitable choices of strings S satisfying (i) and (ii).

Definition 7.3. Let B be a band for Λ . A word S is called a top-socle piece of B if

- a. $S \in \{\alpha^{-1}, \beta^{-1}, \gamma^{-1}, \eta^{-1}, \alpha^{-1}\beta^{-1}, \beta^{-1}\gamma^{-1}, \gamma^{-1}\alpha^{-1}\}$, and
- b. $B \sim_r ST$ for some string T where $T = \xi T' \zeta$ and ξ, ζ are arrows in Q .

Note that $B \sim_r C_0C_1^{-1}C_2C_3^{-1} \dots C_s^{-1}$ where $s \geq 1$ is odd and $C_0, C_1, \dots, C_s$ are top-socle pieces of B .

In the proof of the following proposition, we will often use the equality sign instead of the more precise $\sim_r$.

Proposition 7.4. Let $\Lambda = kQ/I$ where Q and I are as in §3. Then each module M which lies in a component of the stable Auslander-Reiten quiver of Λ which is a 1-tube satisfies $\dim_k \underline{\text{End}}_\Lambda(M) \geq 2$.

Proof. Consider first the 1-tube consisting of string modules. If X is a string module in this 1-tube, then $X = M(\beta^{-1}\gamma^{-1}(\alpha\beta^{-1}\gamma^{-1})^n)$ for some $n \geq 0$. Let $\{z_r\}_{r=0}^{3n+2}$ be the canonical k -basis of the string module X . Then the endomorphism sending z_0 to z_{3n+2} and all other z_r to zero does not factor through a projective Λ -module.

All other 1-tubes consist of band modules, and it follows from [15] that the only possible band modules with stable endomorphism ring k are those lying at the boundary. Hence one only needs to consider the band modules $M_{B,\lambda} = M(B, \lambda, 1)$ where B is a band for Λ and $\lambda \in k - \{0\}$. Suppose there is a band B such that $M_{B,\lambda}$ has stable endomorphism ring equal to k .

Since every band contains either $\beta^{-1}\gamma^{-1}$ or η^{-1} as top-socle piece, one successively eliminates these two cases by determining endomorphisms of suitable string types which do not factor through a projective Λ -module. We provide a few examples in each case.

Suppose first that B contains $\beta^{-1}\gamma^{-1}$ as top-socle piece, i.e. $B = \beta^{-1}\gamma^{-1}C$ for some string $C = C_1^{-1}C_2C_3^{-1} \dots C_s^{-1}$ where $s \geq 1$ is odd and $C_1, C_2, \dots, C_s$ are top-socle pieces of B . One needs to distinguish between $s = 1$ and $s \geq 3$. If $s \geq 3$ then, without loss of generality, $C_1^{-1}C_2 \neq \alpha\beta^{-1}\gamma^{-1}$.

This leads to two cases, namely either $C_1^{-1}C_2 = \alpha\gamma\eta^{-1}$ or $C_1^{-1}C_2 = \alpha\beta^{-1}$. For example, if $C_1^{-1}C_2 = \alpha\gamma\eta^{-1}$ then $M_{B,\lambda}$ has an endomorphism of string type γ or of string type $\gamma^{-1}\alpha$ which does not factor through a projective Λ -module.

Suppose now that B contains η^{-1} as top-socle piece but not $\beta^{-1}\gamma^{-1}$. One needs to distinguish between the case when $B = \eta^{-1}\beta\alpha^{-1}\gamma$ and the case when B also contains $\alpha^{-1}\beta^{-1}$ as top-socle piece. If B contains both η^{-1} and $\alpha^{-1}\beta^{-1}$ as top-socle pieces then $B = \eta^{-1}\beta\alpha^{-1}\beta^{-1}\eta D$ for some string $D = D_1D_2^{-1}D_3 \cdots D_u^{-1}$ where $u \geq 2$ is even and $D_1, D_2, \dots, D_u$ are top-socle pieces of B . This leads to two subcases $D_1 = \gamma^{-1}\alpha^{-1}$ or $D_1 = \gamma^{-1}$. Each of these subcases leads to further subcases that have to be distinguished. For example, if $D_1 = \gamma^{-1}$, then either $D_2^{-1} = \alpha$ or $D_2^{-1} = \alpha\gamma$. Concentrating on the case when $D_1D_2^{-1} = \gamma^{-1}\alpha$, one has $u \geq 4$ and, by our assumptions, $D_3D_4^{-1} = \beta^{-1}\eta$. Then $M_{B,\lambda}$ has an endomorphism of string type $\beta^{-1}\eta\gamma^{-1}$ or of string type $\alpha\beta^{-1}\eta\gamma^{-1}\alpha$ which does not factor through a projective Λ -module.

Altogether, it follows that B contains neither $\beta^{-1}\gamma^{-1}$ nor η^{-1} as top-socle piece, which means that B does not exist and hence Proposition 7.4 follows. $\square$

REFERENCES

- [1] J. L. Alperin, Local representation theory. Modular representations as an introduction to the local representation theory of finite groups. Cambridge Studies in Advanced Mathematics, vol. 11, Cambridge University Press, Cambridge, 1986.
- [2] M. Auslander and J. F. Carlson, Almost-split sequences and group rings. *J. Algebra* 103 (1986), 122–140.
- [3] F. M. Bleher, Universal deformation rings and Klein four defect groups. *Trans. Amer. Math. Soc.* 354 (2002), 3893–3906.
- [4] F. M. Bleher, Universal deformation rings and dihedral defect groups. In press, *Trans. Amer. Math. Soc.*.
- [5] F. M. Bleher, Universal deformation rings for dihedral 2-groups. In press, *J. London Math. Soc.*.
- [6] F. M. Bleher and T. Chinburg, Universal deformation rings and cyclic blocks. *Math. Ann.* 318 (2000), 805–836.
- [7] F. M. Bleher and T. Chinburg, Universal deformation rings need not be complete intersections. *C. R. Math. Acad. Sci. Paris* 342 (2006), 229–232.
- [8] F. M. Bleher and T. Chinburg, Universal deformation rings need not be complete intersections. *Math. Ann.* 337 (2007), 739–767.
- [9] C. Breuil, B. Conrad, F. Diamond and R. Taylor, On the modularity of elliptic curves over $\mathbb{Q}$: Wild 3-adic exercises. *J. Amer. Math. Soc.* 14 (2001), 843–939.
- [10] M. C. R. Butler and C. M. Ringel, Auslander-Reiten sequences with few middle terms and applications to string algebras. *Comm. Algebra* 15 (1987), 145–179.
- [11] J. F. Carlson and J. Thévenaz, The classification of endo-trivial modules. *Invent. Math.* 158 (2004), 389–411.
- [12] G. Cornell, J. H. Silverman and G. Stevens (eds.), *Modular Forms and Fermat’s Last Theorem* (Boston, 1995). Springer-Verlag, Berlin-Heidelberg-New York, 1997.
- [13] B. de Smit and H. W. Lenstra, Explicit construction of universal deformation rings. In: *Modular Forms and Fermat’s Last Theorem* (Boston, MA, 1995), Springer-Verlag, Berlin-Heidelberg-New York, 1997, pp. 313–326.
- [14] K. Erdmann, Blocks of Tame Representation Type and Related Algebras, *Lecture Notes in Mathematics*, vol. 1428, Springer-Verlag, Berlin-Heidelberg-New York, 1990.
- [15] H. Krause, Maps between tree and band modules. *J. Algebra* 137 (1991), 186–194.
- [16] M. Linckelmann, A derived equivalence for blocks with dihedral defect groups. *J. Algebra* 164 (1994), 244–255.
- [17] M. Linckelmann, The source algebras of blocks with a Klein four defect group, *J. Algebra* 167 (1994), 821–854.
- [18] G. Llorent. Stable endomorphism rings and Ext groups for the symmetric group S_4 . Dissertation, University of Iowa, 2007.
- [19] B. Mazur, Deforming Galois representations. In: *Galois groups over $\mathbb{Q}$* (Berkeley, CA, 1987), Springer-Verlag, Berlin-Heidelberg-New York, 1989, pp. 385–437.
- [20] R. Taylor and A. Wiles, Ring-theoretic properties of certain Hecke algebras. *Ann. of Math.* 141 (1995), 553–572.
- [21] A. Wiles, Modular elliptic curves and Fermat’s last theorem. *Ann. of Math.* 141 (1995), 443–551.

F.B.: DEPARTMENT OF MATHEMATICS, UNIVERSITY OF IOWA, IOWA CITY, IA 52242-1419, U.S.A.
E-mail address: fbleher@math.uiowa.edu

G.L.: DEPARTMENT OF MATHEMATICS, CSU SAN BERNARDINO, CA 92407-2397, U.S.A.
E-mail address: gllorent@csusb.edu